

GEOLOGICAL SURVEY OF WYOMING

University of Wyoming

Laramie, Wyoming

INFORMATION CIRCULAR

February, 1945

W Y O M I N G J A D E

Jade was discovered along the Sweetwater River in southeastern Fremont County, Wyoming, in about 1936, but it has only been within the past four years that the mineral has attracted great attention as a gem stone.

Mineralogy of jade - The name jade is applied to two distinctly different minerals, both of which have somewhat similar colors and physical features, such as hardness and toughness. Both will take a high polish to produce gem stones or ornamental stones.

The mineral jadeite is a member of the pyroxene group of minerals and is a sodium-aluminum silicate. It is hard and tough and the color varies from white to greenish-white to emerald green. The Mogoung district of Upper Burma has been an important source for jadeite, but it is also found in Yunnan in southern China, in Tibet, Mexico, and South America. No jadeite has been definitely identified in Wyoming.

The second variety of jade is the mineral nephrite, and Wyoming jade is of this sort. Nephrite belongs to the amphibole group of minerals and is a lime-magnesium-aluminum silicate. Nephrite has a hardness of about 6.5 on the Mohs scale, that is, it is not quite as hard as quartz or agate. Nephrite is also very tough. The specific gravity

is about 3.0. The color varies from leaf-green to dark-green to black and the mineral may be translucent or opaque. In addition to the Wyoming occurrence, nephrite has been found in parts of China, in Turkestan, Siberia, New Zealand, and Alaska.

Wyoming nephrite varies in color from light-green and translucent, to dark-green, to black and opaque. In addition to the color variations, there are differences in quality in that not all specimens will take a high polish. Frequently the jade contains small dark specks which are softer and leave pits in the polished surfaces. A considerable amount of the jade found is of inferior quality because of this feature.

The light-green translucent nephrite is most desirable for gemstones. The average price for good quality uncut stone of this color is about \$3.00 per pound. The highest price is about \$5.00 per pound, although small lots or individual pieces have been sold at a higher price. The dark-green and black varieties are not as desirable and command much lower prices.

Geological occurrence - The Sweetwater River flows from the southern end of the Wind River Mountains across a terrain made up of pre-Cambrian granite hills protruding from a mantle of Tertiary sediments. The jade occurs (1) in place in pre-Cambrian rocks and (2) as boulders in alluvial deposits. The jade originated in the pre-Cambrian rocks and was eroded and transported by streams to form boulders in the Tertiary beds. Later erosion of the Tertiary conglomerates brought about the distribution of jade boulders along the present stream valleys. Jade also occurs as residual boulders on the inter-stream divides.

Five jade deposits are known in the pre-Cambrian rocks, where jade is associated with granite and ultrabasic rocks. All these deposits are dark-green or black nephrite.

Float boulders of dark-green or black jade have been found over a rather large area along the Sweetwater River between Crook's Creek on the east and the Wind River Mountains on the west. The average size of the boulders is from 6 inches to 1 foot in diameter, although many large boulders have been found.

Boulders of the light-green translucent variety of jade have been found in abundance in only two relatively limited areas. The largest boulder of this type of jade so far reported weighed about 3,200 pounds. On the basis of the general geological structure it appears that the original pre-Cambrian source of the light-green jade may have been buried by later sediments and thus may never be located.

Land- status - The area over which the jade occurs comprises private and federal land, and a small amount of state land. A number of placer claims have been filed, especially in the area where the light-green jade occurs. Lode claims have been filed on three of the deposits of dark-green jade where it occurs in place.

Future possibilities - Gem quality jade occurs essentially in only two restricted areas in the Sweetwater district, comprising about 15 square miles. These areas have been so thoroughly picked over during the past three years that today an experienced jade hunter may search for several days without finding specimens of value. Even when polishing tests must be made to determine the quality of the jade, and a considerable part is of inferior quality.

In addition, there are two other green rocks which occur commonly in the area and which superficially resemble jade. The abundance of these rocks, coupled with the fact that they are so often mistaken for jade, has given rise to exaggerated stories of the abundance of jade in the area. Truckloads of green serpentine have been transported for miles only to be found worthless. A green quartzite is also found in the area and has frequently been confused with jade.

It is possible, of course, that new areas containing gem quality jade will be found, but the entire region has been fairly well combed by present prospectors. None of the jade collectors has depended entirely on collecting and selling jade as a livelihood for any length of time. The average jade hunter has simply found it to be a pleasant way to spend spare time.

In addition to its use as a gem stone the nephrite holds possibilities for industrial uses, and investigations are being made of its potential uses as an industrial mineral.

Lapidarists and collectors - The towns of Lander and Riverton are the centers of jade collecting and lapidary work. Persons interested in purchasing uncut jade, jade jewelry, or art objects may obtain the names of reputable dealers by writing to the Chamber of Commerce of either town.