

EXPLANATION

CORRELATION OF MAP UNITS

Qa	Qac	Qol	Qt	Holocene	QUATERNARY	
Qol						Pleistocene
Qol						
UNCONFORMITY						
To				Miocene	TERTIARY	
Ta						
UNCONFORMITY						
Twr				Oligocene		

DESCRIPTION OF MAP UNITS


HOLOCENE AND PLEISTOCENE SURFICIAL DEPOSITS

- Qa** Alluvial deposits - Unconsolidated and poorly consolidated clay, silt, sand, and gravel mainly in floodplains and lowest terraces. Thickness 0 to 85 feet (Lowry and Crist, 1967).
- Qac** Mixed alluvium and colluvium - Sand, silt, clay, and gravel deposited mainly along intermittent streams; includes slope wash and smaller alluvial fan deposits that coalesce with alluvium. Thickness 0 to approximately 60 feet.
- Qol** Windblown sand and loess - Active and stabilized dunes, composed of loess and very fine-grained sand. Two areas are indicated in the map, one north of Pine Bluffs and the other north of Horse Creek in the northwestern part of the map. These features commonly are located downwind from deflation depressions. Numerous small-scale examples of these types of features occur southeast, east, and northeast of Cheyenne and are mapped with the Ogallala Formation. Thickness 0 to 60 feet.
- Qt** Terrace deposits - Beds of coarse sand and gravel with some boulders and lenses of silt and clay. Includes fragments of weathered granite and other igneous rocks. These terraces occur along present drainages, a few feet to over 100 feet above modern floodplains. Thickness 0 to 65 feet along Horse Creek (Lowry and Crist, 1967).
- Qol** Older alluvial fan deposits (Pleistocene) - Poorly sorted clay, silt, sand, and gravel crudely bedded to nonbedded. Located in the southeastern part of the map above present drainages; currently inactive. Thickness 0 to approximately 150 feet (Rapp and others, 1953).
- Qol** Older terrace deposits (Pleistocene) - Beds of coarse sand and gravel with some boulders and thin lenses of silt and clay. A large portion of the sand and gravel is made up of weathered granite fragments transported from the Laramie Mountains or eroded from the underlying Ogallala and Arkkaree Formations. These are upper level terraces which were deposited in channels cut into the Ogallala and Arkkaree Formations. May represent segments of a dissected older alluvial fan complex in some areas. Thickness 0 to 150 feet (Rapp and others, 1953).
- To** OGALLALA FORMATION (UPPER AND MIDDLE MIOCENE) - Tan to brown, unconsolidated to well-cemented sandstone, siltstone, volcanic ash, and conglomerate (gravel- to boulder-sized), interbedded with claystone and thin beds of limestone toward the east. The conglomerate clasts are primarily Precambrian igneous material, with limestone clasts predominating in some conglomerates near the base of the formation. The deposits are mostly coarse grained near the Laramie Mountains and become finer to the east. Thin, light carbonate-cemented lenses of gravel, sand, and silt, locally referred to as "mortar beds", occur in the lower part of the formation in the central and eastern portions of the map area. Reddish brown gravels of possible Pliocene age are mapped with the Ogallala Formation in areas north, east, and southeast of Cheyenne (Cooley, 1991). Thickness 0 to 330 feet (Lowry and Crist, 1967).
- Ta** ARKKAREE FORMATION (LOWER MIOCENE AND UPPER OLIGOCENE) - Friable, gray, tuffaceous, calcareous, fine-grained, crossbedded sandstone, interbedded with lenses of siltstone and volcanic ash. Elongate, cylindrical, calcareous concretions, 2 to 3 feet thick and up to 10 feet long, occur locally in the sandstones. These concretions have a predominant northwest-southeast alignment (Rapp and others, 1953). Coarse conglomerate occurs locally near the base of the formation. The Arkaree Formation pinches out toward the south on the map. Thickness 0 to 500 feet (Lowry and Crist, 1967).
- Twr** WHITE RIVER FORMATION (OLIGOCENE) - The upper White River Formation is characterized by light pink to buff, massive, argillaceous, calcareous siltstone interbedded with minor sandstone, conglomerate, and volcanic ash beds. This unit commonly weathers into cubic blocks and slabs. The silt is imbedded in a matrix of carbonate flakes, volcanic ash, and montmorillonitic clay. The lower White River Formation consists of interbedded red and green bentonitic claystone, sandstone, siltstone, and conglomerate. This lower unit does not crop out in the map area. In adjacent areas, the White River is subdivided into an upper Bruie Formation and a lower Chadron Formation corresponding to the upper and lower White River Formation described above. Thickness 0 to 900 feet (Lowry and Crist, 1967).

SYMBOLS

- CONTACT** - Dashed where approximately located.
- ANTICLINE** - Trace of axis and direction of plunge. Shorter arrow indicates limb of fold with steeper dip. Dashed where approximately located. Plotted axes are based on structure contours on the White River Formation from a map by Lowry and Crist (1967). Axis in Cheyenne area portrays a near-surface fold in the White River Formation.

LOCATION DIAGRAM



600 copies printed by Pikes Peak Lithographing Co., Colorado Springs, Colorado.

Scale 1:100,000

1 CENTIMETER ON THE MAP REPRESENTS 100 METERS ON THE GROUND

CONTOUR INTERVAL 20 METERS

Scale bars in meters and miles.

GEOLOGIC MAP OF THE CHEYENNE 30' x 60' QUADRANGLE, SOUTHEASTERN WYOMING, WESTERN NEBRASKA, AND NORTHERN COLORADO

compiled and mapped by
Alan J. Ver Ploeg
1995

People with disabilities who require an alternate form of communication in order to use this publication should contact the Editor, Wyoming State Geological Survey, TDD Relay operator 1 (800) 877-9975.

Printed on 50% recycled fiber paper.

NORTH

UTM grid convergence (Qa) and 1981 magnetic declination (M) at center of map. Diagram is approximate.

GEOLOGIC SOURCES

Number corresponds to mapped area; unnumbered items are general references.

Babcock, H.M., and Bjorkland, L.J., 1956, Ground-water geology of parts of Laramie and Albany Counties, Wyoming, and Weld County, Colorado. U.S. Geological Survey Water-Supply Paper 1387, 61 p.

1. Bjorkland, L.J., 1957, Geology and ground water resources of the upper Lodgepole Creek drainage basin, Nebraska. U.S. Geological Survey Water-Supply Paper 1410, 76 p., plate 1, scale 1:125,000.

2. Bjorkland, L.J., 1959, Geology and ground water resources of the upper Lodgepole Creek drainage basin, Wyoming. U.S. Geological Survey Water-Supply Paper 1483, 40 p., plate 1, scale 1:126,720.

3. Burchett, R.R., 1986, Geologic bedrock map of Nebraska: Nebraska Geological Survey, 1 plate, scale 1:1,000,000.

4. Cassiliano, Michael, 1980, Stratigraphy and vertebrate paleontology of the Horse Creek-Trail Creek area, Laramie County, Wyoming. University of Wyoming Contributions to Geology, v. 19, n. 1, p. 25-68, figure 3, scale 1:100,000.

5. Cooley, M.E., 1991, Quaternary geology of the northern Great Plains—The Cheyenne tableland, in Morrison, R.B., editor, Quaternary nonglacial geology, continental U.S.: Geological Society of America, The Geology of North America, D.N.A.G. v. K-2, p. 453-455, figure 10, scale 1:400,000.

Cooley, M.E., and Crist, M.A., 1981, Generalized fence diagram showing stratigraphy and potentiometric surfaces of the Tertiary formations in southeast Wyoming and adjacent parts of Colorado. U.S. Geological Survey Miscellaneous Investigations Series Map I-1308, vertical scale 1 inch = 800 feet.

6. Cooley, M.E., and Crist, M.A., 1994, Geohydrology of the High Plains aquifer system, Cheyenne urban area, Wyoming. U.S. Geological Survey Water Resources Investigations Report 92-40477, sheet 3, scale 1:15,840.

Darton, N.H., 1903, Preliminary report on the geology and water resources of Nebraska east of the one hundred and third meridian. U.S. Geological Survey Professional Paper 17, 69 p.

Darton, N.H., 1905, Preliminary report on geology and underground water resources of the central Great Plains. U.S. Geological Survey Professional Paper 32, 433 p.

Denson, N.M., and Bergendahl, M.H., 1961, Middle and upper Tertiary rocks of southeastern Wyoming and adjoining areas. U.S. Geological Survey Professional Paper 424-C, p. C168-C172.

Flanagan, K.L., and Montagne, John, 1993, Neogene stratigraphy and tectonics of Wyoming, in Snoke, A.W., Steidtmann, J.R., and Roberts, S.M., editors, Geology of Wyoming: Wyoming State Geological Survey Memoir 5, p. 572-607.

7. Kirkham, R.M., and Weld, J.W., 1986, Water resources of upper Crow Creek, Colorado. Colorado Geological Survey Special Publication 29, 102 p., plate 1, scale 1:100,000.

Love, J.D., and Christiansen, A.C., 1985, Geologic map of Wyoming. U.S. Geological Survey 3 plates, scale 1:500,000.

Lowry, M.E., 1966, The White River Formation as an aquifer in southeastern Wyoming and adjacent parts of Nebraska and Colorado. U.S. Geological Survey Professional Paper 550-D, p. D217-D222.

8. Lowry, M.E., and Crist, M.A., 1967, Geology and ground water resources of Laramie County, Wyoming. U.S. Geological Survey Water-Supply Paper 1834, 71 p., plate 1, scale 1:125,000.

9. Moore, F.E., 1959, The geomorphic evolution of the east flank of the Laramie Range [sic], Colorado and Wyoming. Laramie, University of Wyoming, unpublished Ph.D. dissertation, 123 p., plate 4, scale 1:63,360.

Moore, F.E., 1963, Tertiary stratigraphy of the High Plains, in Katich, P.J., and Bolyard, D.W., editors, Guidebook to the geology of the northern Denver Basin and adjacent uplifts: Rocky Mountain Association of Geologists, p. 162-166.

10. Rapp, J.R., Warner, D.A., and Morgan, A.M., 1953, Geology and water resources of the Egbert-Pine Bluffs-Carpenter area, Laramie County, Wyoming. U.S. Geological Survey Water-Supply Paper 1140, 67 p., plate 1, scale 1:90,000.

11. State of Wyoming, 1987, Site proposal—superconducting super collider (SSC) U.S. Department of Energy-Geology and tunneling: State of Wyoming, v. 3, 57 p., figure 3.2-1-1a, scale 1:24,000.

Stevenson, Abe, Lloyd, M.D., and Joseph, Lionel, 1983, Soil survey of Laramie County, eastern part: U.S. Department of Agriculture Soil Conservation Service, Washington, D.C., 117 p.

Tweto, Ogden, 1979, Geologic map of Colorado. U.S. Geological Survey, sheet 1, scale 1:500,000.

Weist, W.G., Jr., 1965, Reconnaissance of the ground-water resources in parts of Larimer, Logan, Morgan, Sedgewick, and Weld Counties, Colorado. U.S. Geological Survey Water-Supply Paper 1809-L, 24 p., plate 1, scale 1:250,000.

INDEX TO GEOLOGIC SOURCES

